

Resource Summary Report

Generated by [NIF](#) on Jul 31, 2026

Goat anti-Mouse IgG (H+L) Secondary Antibody, DyLight™ 550

RRID:AB_10942171

Type: Antibody

Proper Citation

(Thermo Fisher Scientific Cat# 84540, RRID:AB_10942171)

Antibody Information

URL: http://antibodyregistry.org/AB_10942171

Proper Citation: (Thermo Fisher Scientific Cat# 84540, RRID:AB_10942171)

Target Antigen: Mouse IgG (H+L)

Host Organism: goat

Clonality: polyclonal secondary

Comments: Applications: Flow, ICC/IF, IHC, IP, WB

Antibody Name: Goat anti-Mouse IgG (H+L) Secondary Antibody, DyLight™ 550

Description: This polyclonal secondary targets Mouse IgG (H+L)

Target Organism: mouse

Defining Citation: [PMID:26141948](#), [PMID:26240283](#)

Antibody ID: AB_10942171

Vendor: Thermo Fisher Scientific

Catalog Number: 84540

Record Creation Time: 20251213T073724+0000

Record Last Update: 20260225T230859+0000

Ratings and Alerts

No rating or validation information has been found for Goat anti-Mouse IgG (H+L) Secondary Antibody, DyLight™ 550.

No alerts have been found for Goat anti-Mouse IgG (H+L) Secondary Antibody, DyLight™ 550.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We found 5 mentions in open access literature.

Listed below are recent publications. The full list is available at [NIF](#).

Brock K, et al. (2024) A comparative analysis of paxillin and Hic-5 proximity interactomes. Cytoskeleton (Hoboken, N.J.).

Papini C, et al. (2022) Exploring APOBEC3A and APOBEC3B substrate specificity and their role in HPV positive head and neck cancer. iScience, 25(10), 105077.

Watanabe S, et al. (2022) Dihydromyricetin improves social isolation-induced cognitive impairments and astrocytic changes in mice. Scientific reports, 12(1), 5899.

Ciancone AM, et al. (2022) Global profiling identifies a stress-responsive tyrosine site on EDC3 regulating biomolecular condensate formation. Cell chemical biology, 29(12), 1709.

Miyoshi T, et al. (2021) Semi-automated single-molecule microscopy screening of fast-dissociating specific antibodies directly from hybridoma cultures. Cell reports, 34(5), 108708.