

Resource Summary Report

Generated by [NIF](#) on Jul 31, 2026

CD11b antibody [MEM-174] (PE-Dyomics647)

RRID:AB_11162994

Type: Antibody

Proper Citation

GeneTex Cat# GTX79991, RRID:AB_11162994

Antibody Information

URL: http://antibodyregistry.org/AB_11162994

Proper Citation: GeneTex Cat# GTX79991, RRID:AB_11162994

Target Antigen: CD11b antibody [MEM-174] (PE-Dyomics647)

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Immunofluorescence; Flow Cytometry; Immunoprecipitation; Flow cytometry, Immunoprecipitation. The usefulness of this product in other applications has not been determined., FACS, IP, Recommended Starting Dilutions: FACS: Positive Control: Peripheral Blood Lymphocytes; IP: Positive Control: GranulocytesCD11b/CD18 transfectants. Note: IP with the antibody MEM-174 was successfully performed using immuno-affinity sorbents (minicolumns). The final immunosorbent should contain 1-5 mg of antibody/1 ml of sorbent gel (e.g. agarose beads) It is recommended to stimulate. Positive Control material by LPS 1-2 days before to achieve better expression of target antigen); Optimal dilutions should be determined experimentally by the researcher.

Antibody Name: CD11b antibody [MEM-174] (PE-Dyomics647)

Description: This monoclonal targets CD11b antibody [MEM-174] (PE-Dyomics647)

Target Organism: human

Antibody ID: AB_11162994

Vendor: GeneTex

Catalog Number: GTX79991

Record Creation Time: 20250819T223529+0000

Ratings and Alerts

No rating or validation information has been found for CD11b antibody [MEM-174] (PE-Dyomics647).

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Immunofluorescence; Flow Cytometry; Immunoprecipitation; Flow cytometry, Immunoprecipitation. The usefulness of this product in other applications has not been determined., FACS, IP, Recommended Starting Dilutions: FACS: Positive Control: Peripheral Blood Lymphocytes; IP: Positive Control: GranulocytesCD11b/CD18 transfectants. Note: IP with the antibody MEM-174 was successfully performed using immuno-affinity sorbents (minicolumns). The final immunosorbent should contain 1-5 mg of antibody/1 ml of sorbent gel (e.g. agarose beads) It is recommended to stimulate. Positive Control material by LPS 1-2 days before to achieve better expression of target antigen); Optimal dilutions should be determined experimentally by the researcher.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.