

Resource Summary Report

Generated by NIF on Jul 31, 2026

c-Myc antibody [9E10] (HRP)

RRID:AB_11165601

Type: Antibody

Proper Citation

GeneTex Cat# GTX80251, RRID:AB_11165601

Antibody Information

URL: http://antibodyregistry.org/AB_11165601

Proper Citation: GeneTex Cat# GTX80251, RRID:AB_11165601

Target Antigen: c-Myc antibody [9E10] (HRP)

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Flow cytometry, Immunohistochemistry, IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western Blot. The usefulness of this product in other applications has not been determined., FACS, IHC, IHC-P, IP, WB, Recommended Starting Dilutions: FACS: Use at 1-5 µg/mL. Note: Membrane permeabilization is required; IP: Use at 1-5 µg/mL. Note: not suitable for IP of native c-Myc protein; WB: Use at 0.5-2 µg/mL. Positive Control: c-Myc tagged protein; IHC-P: Use at 5-10 µg/mL. Positive tissue: perfused brain sections, liver, spleen; Optimal dilutions should be determined experimentally by the researcher.; Immunoprecipitation; Flow Cytometry; Western Blot; Immunofluorescence; Immunohistochemistry; Immunohistochemistry - fixed

Antibody Name: c-Myc antibody [9E10] (HRP)

Description: This monoclonal targets c-Myc antibody [9E10] (HRP)

Target Organism: other, human

Antibody ID: AB_11165601

Vendor: GeneTex

Catalog Number: GTX80251

Record Creation Time: 20250819T223115+0000

Ratings and Alerts

No rating or validation information has been found for c-Myc antibody [9E10] (HRP).

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Flow cytometry, Immunohistochemistry, IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western Blot. The usefulness of this product in other applications has not been determined., FACS, IHC, IHC-P, IP, WB, Recommended Starting Dilutions: FACS: Use at 1-5 µg/mL. Note: Membrane permeabilization is required; IP: Use at 1-5 µg/mL. Note: not suitable for IP of native c-Myc protein; WB: Use at 0.5-2 µg/mL. Positive Control: c-Myc tagged protein; IHC-P: Use at 5-10 µg/mL. Positive tissue: perfused brain sections, liver, spleen; Optimal dilutions should be determined experimentally by the researcher.; Immunoprecipitation; Flow Cytometry; Western Blot; Immunofluorescence; Immunohistochemistry; Immunohistochemistry - fixed

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.