

Resource Summary Report

Generated by [NIF](#) on Jul 28, 2026

Epstein-Barr virus gp 250/350 antibody [0221]

RRID:AB_11165814

Type: Antibody

Proper Citation

GeneTex Cat# GTX36648, RRID:AB_11165814

Antibody Information

URL: http://antibodyregistry.org/AB_11165814

Proper Citation: GeneTex Cat# GTX36648, RRID:AB_11165814

Target Antigen: Epstein-Barr virus gp 250/350 antibody [0221]

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Immunohistochemistry; ELISA; Immunofluorescence; Immunocytochemistry; Other; Western Blot; Recommended Starting Dilutions: For ICC/IF: Use at 1:10-1:50; For WB: Use at 1:10-1:50; For IHC: Use at 1:10-1:50; For ELISA: Use at 1:20-1:200; Not yet tested in other applications. Optimal dilutions should be determined experimentally by the researcher.

For immunostaining cells containing EBNA-1 with clone 0211, slides must be fixed in methanol/acetone (3:7) at -20 °C. After drying at room temperature, slides should be stored at 4°C (1 month) or -20°C (6-12 months). A highly amplified assay must be used to detect the low amount of EBNA expressed, i.e. Monoclonal antibody + Goat Anti-Mouse IgG + Rabbit Anti-Goat IgG-FITC conj., Immunocytochemistry, Immunofluorescence, Western Blot. Th

Antibody Name: Epstein-Barr virus gp 250/350 antibody [0221]

Description: This monoclonal targets Epstein-Barr virus gp 250/350 antibody [0221]

Target Organism: virus

Antibody ID: AB_11165814

Vendor: GeneTex

Catalog Number: GTX36648

Record Creation Time: 20250819T231245+0000

Record Last Update: 20250820T043836+0000

Ratings and Alerts

No rating or validation information has been found for Epstein-Barr virus gp 250/350 antibody [0221].

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Immunohistochemistry; ELISA; Immunofluorescence; Immunocytochemistry; Other; Western Blot; Recommended Starting Dilutions: For ICC/IF: Use at 1:10-1:50; For WB: Use at 1:10-1:50; For IHC: Use at 1:10-1:50; For ELISA: Use at 1:20-1:200; Not yet tested in other applications. Optimal dilutions should be determined experimentally by the researcher.

For immunostaining cells containing EBNA-1 with clone 0211, slides must be fixed in methanol/acetone (3:7) at -20 °C. After drying at room temperature, slides should be stored at 4°C (1 month) or -20°C (6-12 months). A highly amplified assay must be used to detect the low amount of EBNA expressed, i.e. Monoclonal antibody + Goat Anti-Mouse IgG + Rabbit Anti-Goat IgG-FITC conj., Immunocytochemistry, Immunofluorescence, Western Blot. Th

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.