

Resource Summary Report

Generated by [NIF](#) on Jul 30, 2026

PAG antibody [MEM-255] (Dyomics 647)

RRID:AB_11177444

Type: Antibody

Proper Citation

GeneTex Cat# GTX79967, RRID:AB_11177444

Antibody Information

URL: http://antibodyregistry.org/AB_11177444

Proper Citation: GeneTex Cat# GTX79967, RRID:AB_11177444

Target Antigen: PAG antibody [MEM-255] (Dyomics 647)

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Immunohistochemistry - fixed; Immunofluorescence; Flow Cytometry; Immunohistochemistry; Western Blot; Immunocytochemistry; Flow cytometry, Immunohistochemistry, IHC (Formalin-fixed paraffin-embedded sections), Western Blot. The usefulness of this product in other applications has not been determined., FACS, IHC, IHC-P, WB, Recommended Starting Dilutions: FACS: Use at 2 µg/mL. Note:intracellular staining, permeabilization required; WB: Note:Csk binding protein is an ubiquitously expressed 46 kDa transmembrane adaptor protein present in membrane microdomains (rafts), which however migrates on SDS-PAGE gels anomalously as an 80 kDa molecule.; IHC-P: Positive tissue:tonsil, spleen; Optimal dilutions should be determined experimentally by the researcher.

Antibody Name: PAG antibody [MEM-255] (Dyomics 647)

Description: This monoclonal targets PAG antibody [MEM-255] (Dyomics 647)

Target Organism: human

Antibody ID: AB_11177444

Vendor: GeneTex

Catalog Number: GTX79967

Record Creation Time: 20250819T231314+0000

Ratings and Alerts

No rating or validation information has been found for PAG antibody [MEM-255] (Dyomics 647).

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Immunohistochemistry - fixed; Immunofluorescence; Flow Cytometry; Immunohistochemistry; Western Blot; Immunocytochemistry; Flow cytometry, Immunohistochemistry, IHC (Formalin-fixed paraffin-embedded sections), Western Blot. The usefulness of this product in other applications has not been determined., FACS, IHC, IHC-P, WB, Recommended Starting Dilutions: FACS: Use at 2 µg/mL. Note:intracellular staining, permeabilization required; WB: Note:Csk binding protein is an ubiquitously expressed 46 kDa transmembrane adaptor protein present in membrane microdomains (rafts), which however migrates on SDS-PAGE gels anomalously as an 80 kDa molecule.; IHC-P: Positive tissue:tonsil, spleen; Optimal dilutions should be determined experimentally by the researcher.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.