

Resource Summary Report

Generated by [NIF](#) on Jul 31, 2026

proDynorphin antibody

RRID:AB_626032

Type: Antibody

Proper Citation

GeneTex Cat# GTX80766, RRID:AB_626032

Antibody Information

URL: http://antibodyregistry.org/AB_626032

Proper Citation: GeneTex Cat# GTX80766, RRID:AB_626032

Target Antigen: proDynorphin antibody

Host Organism: guinea pig

Clonality: polyclonal

Comments: Discontinued; manufacturer recommendations: Immunohistochemistry 1:500 (Dilutions listed only as a recommendation. Optimal dilution should be determined by investigator.) Antiserum was used on perfusion fixed tissue. Perfusion: 1) calcium-free Tyrode's solution, 2) fixative, and 3) 10 sucrose in PBS as a cryo-protectant. Desired tissues were dissected and stored overnight in 10 sucrose in PBS. Slide-mounted tissue sections were processed for indirect immunofluorescence. Slides were incubated with blocking buffer for 1 hour at room temperature. Primary antiserum was diluted with blocking buffer to the appropriate working concentration. Blocking buffer was removed and slides were incubated for 18-24 hours at 4°C with primary antiserum. Slides were rinsed 3 times and then incubated with secondary antibodies for 1 hour at room temp

Antibody Name: proDynorphin antibody

Description: This polyclonal targets proDynorphin antibody

Target Organism: rat

Antibody ID: AB_626032

Vendor: GeneTex

Catalog Number: GTX80766

Record Creation Time: 20250820T031935+0000

Ratings and Alerts

No rating or validation information has been found for proDynorphin antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Immunohistochemistry 1:500 (Dilutions listed only as a recommendation. Optimal dilution should be determined by investigator.) Antiserum was used on perfusion fixed tissue. Perfusion: 1) calcium-free Tyrode's solution, 2) fixative, and 3) 10 sucrose in PBS as a cryo-protectant. Desired tissues were dissected and stored overnight in 10 sucrose in PBS. Slide-mounted tissue sections were processed for indirect immunofluorescence. Slides were incubated with blocking buffer for 1 hour at room temperature. Primary antiserum was diluted with blocking buffer to the appropriate working concentration. Blocking buffer was removed and slides were incubated for 18-24 hours at 4°C with primary antiserum. Slides were rinsed 3 times and then incubated with secondary antibodies for 1 hour at room temp

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.