

Resource Summary Report

Generated by [NIF](#) on Sep 15, 2026

B6.Cg-E2f1^{Tg(Wnt1-cre)}2Sor/J

RRID:IMSR_JAX:022501

Type: Organism

Proper Citation

RRID:IMSR_JAX:022501

Organism Information

URL: <https://www.jax.org/strain/022501>

Proper Citation: RRID:IMSR_JAX:022501

Description: Mus musculus with name B6.Cg-E2f1^{Tg(Wnt1-cre)}2Sor/J from IMSR.

Species: Mus musculus

Synonyms: B6.Cg-Tg(Wnt1-cre)2Sor/J

Notes: gene symbol note: wingless-type MMTV integration site family; member 1|E2F transcription factor 1|; mutant strain|congenic strain: Wnt1|E2f1|

Affected Gene: wingless-type MMTV integration site family; member 1|E2F transcription factor 1|

Genomic Alteration: transgene insertion 2; Philippe Soriano

Catalog Number: JAX:022501

Database: JAX Mice and Services

Database Abbreviation: JAX

Availability: live

Organism Name: B6.Cg-E2f1^{Tg(Wnt1-cre)}2Sor/J

Record Creation Time: 20250513T053745+0000

Record Last Update: 20260912T060530+0000

Ratings and Alerts

No rating or validation information has been found for B6.Cg-E2f1^{Tg(Wnt1-cre)}2Sor/J.

No alerts have been found for B6.Cg-E2f1^{Tg(Wnt1-cre)}2Sor/J.

Data and Source Information

Source: [Integrated Animals](#)

Source Database: JAX Mice and Services

Usage and Citation Metrics

We found 26 mentions in open access literature.

Listed below are recent publications. The full list is available at [NIF](#) .

Iyyanar PPR, et al. (2026) Cse1l is critical for cell survival, craniofacial and cardiac development. Developmental biology, 530, 188.

Huber J, et al. (2026) The Sh3Pxd2bnee^{-/-} mouse: an attractive model unveiling novel developmental features in Frank-ter-Haar Syndrome. Development (Cambridge, England).

Ma F, et al. (2026) CSF1R⁺ macrophage and osteoclast depletion impairs neural crest proliferation and craniofacial morphogenesis. Development (Cambridge, England), 153(16).

Raymundo JR, et al. (2026) KCTD1/KCTD15 complexes repress AP-2 γ and AP-2 δ to regulate neural crest-dependent craniofacial morphogenesis and scalp skin development. The Journal of investigative dermatology.

Khot B, et al. (2025) miR-219a-Robo1 axis acts as a conserved switch for slit-mediated commissural axon guidance. iScience, 28(12), 114242.

Chakraborty S, et al. (2025) Deletion of a single CTCF motif at the boundary of a chromatin domain with three FGF genes disrupts gene expression and embryonic development. Developmental cell, 60(13), 1838.

Seguella L, et al. (2025) Hyperactive enteric glia contribute to persistent dysmotility following inflammation by driving aberrant excitatory responses in neurons. Cellular and molecular gastroenterology and hepatology, 101634.

Chen P, et al. (2025) Pyroptosis of periodontal ligament stem cells aggravates periodontitis via Piezo1 channel. iScience, 28(5), 112395.

Moore AC, et al. (2025) Role of soft tissue and bone interactions in the developmental integration and modularity of the skull in neural crest-specific gap junction α -1 knockout mice. Anatomical record (Hoboken, N.J. : 2007).

Wong CK, et al. (2024) Central glucagon-like peptide 1 receptor activation inhibits Toll-like receptor agonist-induced inflammation. Cell metabolism, 36(1), 130.

Lehr S, et al. (2024) Self-organized pattern formation in the developing mouse neural tube by a temporal relay of BMP signaling. *Developmental cell*.

Morales-Soto W, et al. (2024) Endocannabinoids regulate enteric neuron-glia networks and visceral hypersensitivity following inflammation through a glial-dependent mechanism. *Glia*.

Hanscom M, et al. (2024) Innervation of adipocytes is limited in mouse perivascular adipose tissue. *American journal of physiology. Heart and circulatory physiology*, 327(1), H155.

Hendler BJ, et al. (2024) Purinergic P2Y1 and P2Y12 receptors control enteric nervous system activity through neuro-glia-macrophage crosstalk. *Purinergic signalling*.

Gunawardhana KL, et al. (2023) A systems biology approach identifies the role of dysregulated PRDM6 in the development of hypertension. *The Journal of clinical investigation*, 133(4).

Raymundo JR, et al. (2023) KCTD1/KCTD15 complexes control ectodermal and neural crest cell functions, and their impairment causes aplasia cutis. *The Journal of clinical investigation*, 134(4).

Ahmadzai MM, et al. (2022) LPAR1 regulates enteric nervous system function through glial signaling and contributes to chronic intestinal pseudo-obstruction. *The Journal of clinical investigation*, 132(4).

Taïb S, et al. (2022) Myelinating Schwann cells and Netrin-1 control intra-nervous vascularization of the developing mouse sciatic nerve. *eLife*, 11.

Van Battum E, et al. (2021) Plexin-B2 controls the timing of differentiation and the motility of cerebellar granule neurons. *eLife*, 10.

Gahankari A, et al. (2021) Deregulated Rac1 Activity in Neural Crest Controls Cell Proliferation, Migration and Differentiation During Midbrain Development. *Frontiers in cell and developmental biology*, 9, 704769.